

Classification of countries of cultivation

The countries¹ of cultivation of maize are classified into 3 risk categories: High, Medium and Low. This classification is carried out on a regular basis by a number of feed schemes together. When determining the risk category of a country of cultivation, the results of aflatoxin analyzes available in the GMP+ Monitoring Database are taken into account, as well as the results collected by other schemes (QS, EFISC-GTP, OVOCOM, AIC). In addition to these results, other relevant information (e.g. RASFF notifications, information about weather/harvest conditions, etc.) can also be used. The criteria used during the evaluation are shown in table 2.

Table 1: Risk classification of countries of cultivation

Publication date: 10 November 2025

Implementation date: 12 November 2025

High	Medium	Low
Brazil Romania Serbia	- All other countries which are not mentioned under 'high risk countries' or 'low risk countries' Southern France (departments: 85, 79, 86, 17, 16, 87, 23, 19, 03, 63, 15, 43, 71, 39, 42, 69, 01, 74, 73, 38, 07, 26, 24, 33, 40, 64, 47, 46, 82, 32, 65, 31, 81, 12, 09, 48, 30, 34, 11, 66, 05, 04, 06, 83, 84 and 13)	Austria Belgium Czech Republic Denmark Estonia Finland France (except Southern France; departments: 85, 79, 86, 17, 16, 87, 23, 19, 03, 63, 15, 43, 71, 39, 42, 69, 01, 74, 73, 38, 07, 26, 24, 33, 40, 64, 47, 46, 82, 32, 65, 31, 81, 12, 09, 48, 30, 34, 11, 66, 05, 04, 06, 83, 84 and 13) Germany Iceland Ireland Latvia Lithuania Luxembourg Netherlands Norway Poland Slovakia Sweden UK

1. If applicable, a country can be divided in different regions.

Table 2: Criteria for (Re)classification of a country of cultivation

Risk classification by country of cultivation	% of analyses per country (period: 1 September until evaluation moment)	Analysis results (x)
High	> 1%	> 20 ppb; OR
	> 10%	10ppb < x ≤ 20ppb
Medium	Any scenario not listed as High or Low	
Low	< 1%	5ppb < x ≤ 10ppb; AND
	> 90%	< 2ppb; AND
	Remaining (max 9%)	≤ 5ppb
a. For upgrading a country of cultivation to a higher risk level, the number of samples to be tested is at least 1. b. For downgrading a country of cultivation to a lower risk level, the number of samples to be tested is at least 50 (new results). c. In addition to a. and b., other relevant criteria and resources can be used for reclassifying a country of origin.		

Previous versions

- [Version 04.09.2025](#)
- [Version 24.04.2025](#)
- [Version 28.03.2025](#)
- [Version 28.02.2025](#)
- [Version 21.11.2024](#)
- [Version 12.09.2024](#)
- [Version 07.06.2024](#)
- [Version 11.04.2024](#)
- [Version 14.03.2024](#)

We enable every company in the feed chain to take responsibility for safe and sustainable feed.

GMP+ International

Braillelaan 9

2289 CL Rijswijk

The Netherlands

t. +31 (0)70 – 307 41 20 (Office)

+31 (0)70 – 307 41 44 (Help Desk)

e. info@gmpplus.org

Disclaimer:

This publication was established for the purpose of providing information to interested parties with respect to GMP+-standards. The publication will be updated regularly. GMP+ International B.V. is not liable for any inaccuracies in this publication.

© GMP+ International B.V.

All rights reserved. The information in this publication may be consulted on the screen, downloaded and printed as long as this is done for your own, non-commercial use. For other desired uses, prior written permission should be obtained from the GMP+ International B.V.